

## Clinical Policy: Talquetamab-tgvs (Talvey)

Reference Number: CP.PHAR.649

Effective Date: 12.01.23

Last Review Date: 11.23

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Talquetamab-tgvs (Talvey<sup>™</sup>) is bispecific GPRC5D-directed CD3 T-cell engager.

### FDA Approved Indication(s)

Talvey is indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antibody.

This indication is approved under accelerated approval based on response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

### Policy/Criteria

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Talvey is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Multiple Myeloma (must meet all):

1. Diagnosis of MM;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq$  18 years;
4. Disease is relapsed or refractory;
5. Member has received or has documented intolerance to  $\geq$  4 prior lines of therapies\* (see Appendix B) that include all of the following (a, b, and c):
  - a. One proteasome inhibitor (e.g., bortezomib, Kyprolis<sup>®</sup>, Ninlaro<sup>®</sup>)
  - b. One immunomodulatory drug (e.g., Thalomid<sup>®</sup>, lenalidomide, pomalidomide)
  - c. One anti-CD38 monoclonal antibodies (e.g., Darzalex<sup>®</sup>, Sarclisa<sup>®</sup>)
6. Request meets one of the following (a, b, or c):\*
  - a. Dose does not exceed 0.4 mg/kg once weekly;
  - b. Dose does not exceed 0.8 mg/kg every 2 weeks;
  - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prior authorization may be required*

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

#### Approval duration:

**Medicaid/HIM** – 6 months

**Commercial** – 6 months or to the member's renewal date, whichever is longer

**B. Other diagnoses/indications** (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**II. Continued Therapy**

**A. Multiple Myeloma** (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Talvey for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a, b, or c):\*
  - a. Dose does not exceed 0.4 mg/kg once weekly;
  - b. Dose does not exceed 0.8 mg/kg every 2 weeks;
  - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration:**

**Medicaid/HIM** – 12 months

**Commercial** – 6 months or to the member’s renewal date, whichever is longer

**B. Other diagnoses/indications** (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

MM: multiple myeloma

NCCN: National Comprehensive Cancer Network

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

<b>Drug Name</b>	<b>Dosing Regimen</b>	<b>Dose Limit/ Maximum Dose</b>
<b>MM: regimens containing proteasome inhibitors, immunomodulatory agents and/or anti-CD38 monoclonal antibodies (examples – NCCN)</b>		
bortezomib / lenalidomide (Revlimid <sup>®</sup> ) or pomalidomide or Thalomid <sup>®</sup> (thalidomide) / dexamethasone	Varies	Varies
Kyprolis <sup>®</sup> (carfilzomib – weekly or twice weekly) / dexamethasone	Varies	Varies
Kyprolis <sup>®</sup> (carfilzomib) / lenalidomide (Revlimid <sup>®</sup> ) / dexamethasone	Varies	Varies
Ninlaro <sup>®</sup> (ixazomib) / lenalidomide (Revlimid <sup>®</sup> ) / dexamethasone	Varies	Varies
Darzalex <sup>®</sup> (daratumumab) / bortezomib / dexamethasone ± Thalomid <sup>®</sup> (thalidomide)	Varies	Varies
Darzalex <sup>®</sup> (daratumumab) / lenalidomide (Revlimid <sup>®</sup> ) / dexamethasone	Varies	Varies

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): None
- Boxed warning(s): cytokine release syndrome, neurologic toxicity

## V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Relapsed or refractory MM	<p>Weekly dosing schedule:</p> <ul style="list-style-type: none"> <li>Day 1: 0.01 mg/kg</li> <li>Day 4: 0.06 mg/kg</li> <li>Day 7 (first treatment dose): 0.4 mg/kg</li> <li>One week after first treatment dose (subsequent treatment doses): 0.4 mg/kg weekly</li> </ul> <p>Biweekly (every 2 weeks) dosing schedule:</p> <ul style="list-style-type: none"> <li>Day 1: 0.01 mg/kg</li> <li>Day 4: 0.06 mg/kg</li> <li>Day 7: 0.4 mg/kg</li> <li>Day 10 (first treatment dose): 0.8 mg/kg</li> <li>Two weeks after first treatment dose (subsequent treatment doses): 0.8 mg/kg every 2 weeks</li> </ul> <p>Dose calculation is based on actual body weight.</p>	0.4 mg/kg once weekly or 0.8 mg/kg every 2 weeks

## VI. Product Availability

Single-dose vials for injection: 3 mg/1.5 mL (2 mg/mL); 40 mg/mL

## VII. References

1. Talvey Prescribing Information. Horsham, PA: Janssen Biotech, Inc.; Aug 2023. Available at: [www.talvey.com](http://www.talvey.com). Accessed August 23, 2023.
2. National Comprehensive Cancer Network. Multiple Myeloma Version 3.2023. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/myeloma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf). Accessed August 23, 2023.
3. Chari A, Minnema MC, Berdeja JG, et al. Talquetamab, a t-cell–redirecting gprc5d bispecific antibody for multiple myeloma. *New England Journal of Medicine*. 2022;387(24):2232-2244.

## Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3055	Injection, talquetamab-tgvs, 0.25 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	08.29.23	11.23
Removed HCPCS codes [C9399, J3590] and added HCPCS code [J3055]	02.20.24	

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members, and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2023 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.